



TITLE:

# Tunneling for spatially cut-off $\phi$ -Hamiltonians

AUTHOR(S):

Aida, Shigeki

---

CITATION:

Aida, Shigeki. Tunneling for spatially cut-off  $\phi$ -Hamiltonians. 数理解析研究所講究録 2013, 1855: 133-139

ISSUE DATE:

2013-10

URL:

<http://hdl.handle.net/2433/195220>

RIGHT:

## Tunneling for spatially cut-off $P(\phi)_2$ -Hamiltonians

Shigeki Aida  
Tohoku University

This note is a short presentation of recent results for semi-classical analysis of lowlying eigenvalues of spatially cut-off  $P(\phi)_2$ -Hamiltonians based on the author's recent research ([2, 3]). We refer the readers for semi-classical analysis in finite dimensions to [17, 21, 32, 33, 19, 20] and for  $P(\phi)_2$ -Hamiltonians to [12, 31, 34, 7].

First, we give a definition of spatially cut-off  $P(\phi)_2$ -Hamiltonians. Let  $m > 0$ . Let  $\mu$  be the Gaussian measure on the space of tempered distributions  $\mathcal{S}'(\mathbb{R})$  such that

$$\int_W \mathcal{S}(\mathbb{R}) \langle \varphi, w \rangle_{\mathcal{S}'(\mathbb{R})}^2 d\mu(w) = \left( (m^2 - \Delta)^{-1/2} \varphi, \varphi \right)_{L^2}.$$

Let  $\mathcal{E}$  be the Dirichlet form defined by

$$\mathcal{E}(f, f) = \int_W \|\nabla f(w)\|_{L^2(\mathbb{R}, dx)}^2 d\mu(w) \quad f \in D(\mathcal{E}),$$

where  $\nabla f(w)$  is the unique element in  $L^2(\mathbb{R}, dx)$  such that

$$\lim_{\varepsilon \rightarrow 0} \frac{f(w + \varepsilon \varphi) - f(w)}{\varepsilon} = (\nabla f(w), \varphi)_{L^2(\mathbb{R}, dx)}.$$

The generator  $-L(\geq 0)$  of  $\mathcal{E}$  is one of expressions of a free Hamiltonian. Let  $P(x) = \sum_{k=0}^{2M} a_k x^k$ , where  $a_{2M} > 0$ . Let  $g \in C_0^\infty(\mathbb{R})$  with  $g(x) \geq 0$  for all  $x$  and define for  $h \in H^1(= H^1(\mathbb{R}))$ ,

$$\begin{aligned} V(h) &= \int_{\mathbb{R}} P(h(x))g(x)dx \\ U(h) &= \frac{1}{4} \int_{\mathbb{R}} (h'(x)^2 + m^2 h(x)^2) dx + V(h) \end{aligned}$$

We want to consider an operator like

$$-L + \lambda V(w/\sqrt{\lambda}) \quad \text{on} \quad L^2(\mathcal{S}'(\mathbb{R}), d\mu).$$

The difficulty is in the definition of  $w(x)^k$  because  $w$  is an element of the Schwartz distribution. Instead of  $w(x)^k$ , we use Wick power  $:w(x)^k:$  which requires renormalizations for which we refer the readers to [12, 31, 34, 7]. For  $P = P(x) = \sum_{k=0}^{2M} a_k x^k$  with  $a_{2M} > 0$ , define

$$\int_{\mathbb{R}} :P\left(\frac{w(x)}{\sqrt{\lambda}}\right): g(x)dx = \sum_{k=0}^{2M} a_k \int_{\mathbb{R}} : \left(\frac{w(x)}{\sqrt{\lambda}}\right)^k : g(x)dx.$$

We write

$$\begin{aligned} : V \left( \frac{w}{\sqrt{\lambda}} \right) &:= \int_{\mathbb{R}} : P \left( \frac{w(x)}{\sqrt{\lambda}} \right) : g(x) dx, \\ V_{\lambda}(w) &= \lambda : V \left( \frac{w}{\sqrt{\lambda}} \right) :. \end{aligned}$$

**Definition 1.** *The spatially cut-off  $P(\phi)_2$ -Hamiltonian  $-L + V_{\lambda}$  is defined to be the unique self-adjoint extension operator of  $(-L + V_{\lambda}, \mathfrak{F}C_b^{\infty}(S'(\mathbb{R})))$ .*

It is known that  $-L + V_{\lambda}$  is bounded from below and the first eigenvalue  $E_1(\lambda)$  is simple and the corresponding positive eigenfunction  $\Omega_{1,\lambda}$  exists. See [12, 31, 34]. Formally,  $-L + V_{\lambda}$  is unitarily equivalent to the infinite dimensional Schrödinger operator:

$$-\Delta_{L^2(\mathbb{R})} + \lambda U(w/\sqrt{\lambda}) - \frac{1}{2} \text{tr}(m^2 - \Delta)^{1/2} \quad \text{on } L^2(L^2(\mathbb{R}), dw)$$

where  $dw$  is an infinite dimensional Lebesgue measure. The function  $U$  is a potential function such that

$$U(w) = \frac{1}{4} \int_{\mathbb{R}} w'(x)^2 dx + \int_{\mathbb{R}} \left( \frac{m^2}{4} w(x)^2 + : P(w(x)) : g(x) \right) dx$$

and  $\Delta_{L^2(\mathbb{R})}$  denotes the “Laplacian” on  $L^2(\mathbb{R}, dx)$ . Hence, by the analogy of Schrödinger operators in finite dimensions, it is natural to expect that asymptotic behavior of lowlying eigenvalues of  $-L + V_{\lambda}$  in the semiclassical limit  $\lambda \rightarrow \infty$  is related with the properties of global minimum points of  $U$ . In view of this, we consider the following assumptions.

**Assumption 2.** *Let  $U$  be the function on  $H^1$  such that*

$$U(h) = \frac{1}{4} \int_{\mathbb{R}} h'(x)^2 dx + \int_{\mathbb{R}} \left( \frac{m^2}{4} h(x)^2 + P(h(x))g(x) \right) dx \quad \text{for } h \in H^1.$$

(A1) *The function  $U$  is non-negative and the zero point set*

$$\mathcal{Z} := \{h \in H^1 \mid U(h) = 0\} = \{h_1, \dots, h_n\}$$

*is a finite set.*

(A2) *For all  $1 \leq i \leq n$ , the Hessian  $\nabla^2 U(h_i)$  is non-degenerate. That is, there exists  $\delta_i > 0$  for each  $i$  such that*

$$\begin{aligned} \nabla^2 U(h_i)(h, h) &:= \frac{1}{2} \int_{\mathbb{R}} h'(x)^2 dx + \int_{\mathbb{R}} \left( \frac{m^2}{2} h(x)^2 + P''(h_i(x))g(x)h(x)^2 \right) dx \\ &\geq \delta_i \|h\|_{L^2(\mathbb{R})}^2 \quad \text{for all } h \in H^1(\mathbb{R}). \end{aligned}$$

(A3) *For all  $x$ ,  $P(x) = P(-x)$  and  $\mathcal{Z} = \{h_0, -h_0\}$ , where  $h_0 \neq 0$ .*

Let  $E_1(\lambda)$  be the lowest eigenvalue of  $-L + V_{\lambda}$ . The first main result is as follows.

**Theorem 3** ([3]). Assume that (A1) and (A2) hold. Let  $E_1(\lambda) = \inf \sigma(-L + V_\lambda)$ . Then

$$\lim_{\lambda \rightarrow \infty} E_1(\lambda) = \min_{1 \leq i \leq n} E_i,$$

where

$$E_i = \inf \sigma(-L + Q_i)$$

and  $Q_i$  is given by

$$Q_i(w) = \frac{1}{2} \int_{\mathbb{R}} : w(x)^2 : P''(h_i(x)) g(x) dx.$$

**Remark 4.** In the case of finite dimensional Schrödinger operators, there exist eigenvalues near the approximate eigenvalues  $E_i$  when  $\lambda$  is large. In Theorem 3, if  $E_i < m + \min_{1 \leq i \leq n} E_i$ , then the same results hold by the result of Hoegh-Krohn and Simon [34]. However, if it is not the case, it is not clear and they may be embedded eigenvalues in the essential spectrum. Under the assumptions in Theorem 6,  $E_2(\lambda)$  is an eigenvalue for large  $\lambda$ . Simon [30] gave an example of  $P(\phi)_2$ -Hamiltonian for which an embedded eigenvalue exists.

Let

$$E_2(\lambda) = \inf \{ \sigma(-L + V_\lambda) \setminus \{E_1(\lambda)\} \}.$$

We can prove that  $E_2(\lambda) - E_1(\lambda)$  is exponentially small when  $U$  is a symmetric double well potential function. The exponential decay rate is given by the Agmon distance which is defined below.

**Definition 5.** Let  $0 < T < \infty$  and  $h, k \in H^1(\mathbb{R})$ . Let  $AC_{T,h,k}(H^1(\mathbb{R}))$  be the set of all absolutely continuous paths  $c : [0, T] \rightarrow H^1(\mathbb{R})$  satisfying  $c(0) = h, c(T) = k$ . Let  $U$  be the potential function in (2). Assume  $U$  is non-negative. We define the Agmon distance between  $h, k$  by

$$d_U^{Ag}(h, k) = \inf \{ \ell_U(c) \mid c \in AC_{T,h,k}(H^1(\mathbb{R})) \},$$

where

$$\ell_U(c) = \int_0^T \sqrt{U(c(t))} \|c'(t)\|_{L^2} dt.$$

The following estimate is the second main result.

**Theorem 6** ([3]). Assume that  $U$  satisfies (A1), (A2), (A3). Then it holds that

$$\limsup_{\lambda \rightarrow \infty} \frac{\log(E_2(\lambda) - E_1(\lambda))}{\lambda} \leq -d_U^{Ag}(h_0, -h_0).$$

**Remark 7.** (1) Agmon distance can be extended to a continuous distance function on  $H^{1/2}(\mathbb{R})$ . Moreover the topology defined by the Agmon distance coincides with the one defined by the Sobolev norm of  $H^{1/2}(\mathbb{R})$ .

(2) We can prove the existence of minimal geodesic between  $h_0$  and  $-h_0$  with respect to the Agmon metric. The uniqueness of the geodesics is not clear at the moment.

(3) The Agmon distance  $d_U^{Ag}(h_0, -h_0)$  is equal to an Euclidean action integral of an instanton solution. This is an infinite dimensional example corresponding to the result of instanton in the case of Schrödinger operator which is due to Carmona and Simon [6].

The following is an example for which our main theorem is applicable.

**Example 8.** Fix  $g \in C_0^\infty(\mathbb{R})$ . Let  $n \in \mathbb{N}$ . For sufficiently large  $a > 0$ , the polynomial

$$P(x) = a(x^2 - 1)^{2n} - C$$

satisfies (A1), (A2), (A3). Here  $C$  is a positive constant which depends on  $a, g$ .

The same theorems are valid in the case where the space is a finite interval  $I = [-l/2, l/2]$  as in the setting in [2]. In that framework, we show a simple example for which the Agmon distance and instanton can be calculated. Let  $a$  and  $x_0$  be positive numbers. We consider the case where

$$U(h) = \frac{1}{4} \int_I h'(x)^2 dx + a \int_I (h(x)^2 - x_0^2)^2 dx.$$

For example, setting  $b^2 = x_0^2 + \frac{m^2}{8a}$  and

$$P(x) = a(x^2 - b^2)^2 - a \left\{ b^4 - \left( b^2 - \frac{m^2}{8a} \right)^2 \right\},$$

we obtain the potential function above. Note  $\mathcal{Z} = \{h_0, -h_0\}$ , where  $h_0(x) \equiv x_0$  is a constant function.  $\pm x_0$  are the zero points also of the potential function

$$Q(x) = a(x^2 - x_0^2)^2 \quad x \in \mathbb{R}.$$

Let

$$d_{1dim}^{Ag}(-x_0, x_0) = \inf \left\{ \int_{-T}^T \sqrt{Q(x(t))} |x'(t)| dt \mid x(-T) = -x_0, \quad x(T) = x_0 \right\}.$$

This is the Agmon distance which corresponds to 1-dimensional Schrödinger operator  $-\frac{d^2}{dx^2} + Q(x)$  defined in  $L^2(\mathbb{R}, dx)$  and

$$d_{1dim}^{Ag}(-x_0, x_0) = \int_{-x_0}^{x_0} \sqrt{Q(x)} dx = \frac{4\sqrt{a}x_0^3}{3}.$$

We can prove the following.

**Proposition 9.** Assume  $2ax_0^2 l^2 \leq \pi^2$ . Let  $u_0(t) = x_0 \tanh(2\sqrt{a}x_0 t)$ . Then  $u_0(t)$  is a solution to

$$\begin{aligned} u''(t) &= 2Q'(u(t)) & -\infty < t < \infty, \\ \lim_{t \rightarrow -\infty} u(t) &= -x_0, & \lim_{t \rightarrow \infty} u(t) = x_0 \end{aligned}$$

and

$$\begin{aligned} I_{\infty, P}(u_0) &= \left( \frac{1}{4} \int_{-\infty}^{\infty} u_0'(t)^2 dt + \int_{-\infty}^{\infty} Q(u_0(t)) dt \right) l, \\ &= d_{1dim}^{Ag}(-x_0, x_0) l \\ &= d_U^{Ag}(-h_0, h_0). \end{aligned}$$

The Proposition above claims that  $u_0$  is the instanton for both operators: 1-dimensional Schrödinger operator  $-\frac{d^2}{dx^2} + \lambda Q(\cdot/\sqrt{\lambda})$  and  $-L + V_\lambda$ .

## References

- [1] S. Agmon, Lectures on Exponential decay of solutions of second order elliptic equations. Bounds on eigenfunctions of  $N$ -body Schrödinger operators, Mathematical Notes, Princeton Univ. Press, Princeton, N.J., 1982.
- [2] S. Aida, Semi-classical limit of the lowest eigenvalue of a Schrödinger operator on a Wiener space : II.  $P(\phi)_2$ -model on a finite volume, J. Funct. Anal. 256 (2009), no. 10, 3342–3367.
- [3] S. Aida, Tunneling for spatially cut-off  $P(\phi)_2$ -Hamiltonians. J. Funct. Anal. 263 (2012), no. 9, 2689—2753.
- [4] A. Arai, Trace formulas, a Golden-Thompson inequality and classical limit in boson Fock space, J. Funct. Anal. **136**, (1996), 510–547.
- [5] C. Borell, Tail probabilities in Gauss space. Vector space measures and applications. I, Lecture Notes in Math. **644**, 71–82 (1978), Springer.
- [6] R. Carmona and B. Simon, Pointwise bounds on eigenfunctions and wave packets in  $N$ -body quantum systems, V. Lower bounds and path integrals, Comm. Math. Phys. **80** (1981), 59–98.
- [7] J. Dereziński and C. Gérard, Spectral scattering theory of spatially cut-off  $P(\phi)_2$  Hamiltonians, Commun. Math. Phys. **213**, 39–125, (2000).
- [8] M. Dimassi and J. Sjöstrand, Spectral asymptotics in the semi-classical limit, London Mathematical Society Lecture Note Series, **268**, Cambridge University Press, 1999.
- [9] S. Y. Dobrokhoto and V. N. Kolokol'tsov, The double-well splitting of low energy levels for the Schrödinger operator of discrete  $\phi^4$ -models on tori, J. Math. Phys. **36** (3), 1038–1053, (1995).
- [10] W.J. Eachus and L. Streit, Exact solution of the quadratic interaction Hamiltonian, Reports on Mathematical Physics **4**, No. 3, 161–182, 1973.
- [11] J. P. Eckmann, Remarks on the classical limit of quantum field theories, Lett. Math. Phys. **1** (1975/1977), no. 5, 387–394.
- [12] 江沢洋・新井朝雄, 場の量子論と統計力学, 日本評論社 (1988).
- [13] P. Federbush, Partially Alternate Derivation of a Result of Nelson, J. Math. Phys. Vol. 10, No. 1 January, 1969
- [14] J. Glimm, Boson fields with non-linear self-interaction in two dimensions, Comm. Math. Phys. **8** (1968), 12–25.
- [15] L. Gross, Abstract Wiener spaces. 1967 Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 1 pp. 31–42 Univ. California Press, Berkeley, Calif.

- [16] L. Gross, Logarithmic Sobolev inequalities, *Amer. J. Math.* **97** (1975), 1061-1083.
- [17] E. Harrell, Double wells. *Comm. Math. Phys.* 75 (1980), no. 3, 239—261.
- [18] B. Helffer and F. Nier, Hypoelliptic estimates and spectral theory for Fokker-Planck operators and Witten Laplacians, *Lecture Notes in Mathematics*, **1862**, Springer, 2005.
- [19] B. Helffer and J. Sjöstrand, Multiple wells in the semiclassical limit. I. *Comm. Partial Differential Equation* **9** (1984), no.4, 337–408.
- [20] B. Helffer and J. Sjöstrand, Puits multiples en limite semi-classique. II. Interaction moléculaire. Symétries. Perturbation. *Ann. Inst. H. Poincaré Phys. Théor.* **42**, no. 2 (1985), 127–212.
- [21] G. Jona-Lasinio, F. Martinelli and E. Scoppola, New approach to the semiclassical limit of quantum mechanics. I. Multiple tunnelings in one dimension. *Comm. Math. Phys.* 80 (1981), no. 2, 223—254.
- [22] H-H, Kuo, Gaussian measures in Banach spaces. *Lecture Notes in Mathematics*, **463**, Springer-Verlag, Berlin-New York, 1975.
- [23] M. Ledoux, Isoperimetry and Gaussian analysis, *Lectures on probability theory and statistics*, (Saint-Flour, 1994), 165–294, *Lecture Notes in Math.*, **1648**, Springer, Berlin, 1996.
- [24] M. Ledoux, A note on large deviations for Wiener chaos, *Séminaire de Probabilités, XXXIV*, 1988/89, 1–14. *Lecture Notes in Math.*, **1426**, Springer, Berlin, 1990.
- [25] L. Lithner, A theorem of the Phragmén-Lindelöf type for second-order elliptic operators. *Ark. Mat.* 5 1964 281—285 (1964).
- [26] O. Matte and J. S. Møller, On the spectrum of semi-classical Witten-Laplacians and Schrödinger operators in large dimension, *J. Funct. Anal.*, **220** (2005), no. 2, 243–264.
- [27] E. Nelson, A quartic interaction in two dimensions. 1966 *Mathematical Theory of Elementary Particles* (Proc. Conf., Dedham, Mass., 1965) 69–73 M.I.T. Press, Cambridge, Mass.
- [28] L. Rosen, Renormalization of the Hilbert space in the mass shift model, *J. Mathematical Phys.* **13** (1972), 918–927.
- [29] I. Segal, Construction of nonlinear local quantum processes. I, *Ann. Math. (2)*, **92** (1970), 462–481.
- [30] B. Simon, Continuum embedded eigenvalues in a spatially cutoff  $P(\phi)_2$  field theory, *Proc. Amer. Math. Soc.* **35** (1972), 223–226.
- [31] B. Simon, *The  $P(\phi)_2$  Euclidean (quantum) field theory*, Princeton University Press, New Jersey, (1974).

- [32] B. Simon, Semiclassical Analysis of Low Lying Eigenvalues I. Nondegenerate Minima: Asymptotic Expansions, *Ann. Inst. Henri Poincaré, Section A*, Vol. XXXVIII, no. 4, (1983), 295–308.
- [33] B. Simon, Semiclassical analysis of low lying eigenvalues, II. Tunneling, *Annals of Math.* **120**, (1984), 89–118.
- [34] B. Simon and R. Hoegh-Krohn, Hypercontractive Semigroups and Two Dimensional Self-Coupled Bose Fields, *J. Funct. Anal.*, Vol. 9, (1972), 121–180.

Mathematical Institute

Tohoku University

Sendai 980-8578

JAPAN

E-mail address: aida@math.tohoku.ac.jp